

THERMOMAGNETIC WAVES IN UNSTABLE ANISOTROPIC CONDUCTING MEDIA

**E.R. HASANOV^{1,2}, Sh.G. KHALILOVA², G.M. MAMMADOVA²,
R.Z. JAVADOVA¹**

¹*Baku State University Acad. Z.Khalilov, str.23, Baku, Azerbaijan Republic*
²*Institute of Physics, ANAS, AZ-1143, H. Javid 131, Baku, Azerbaijan Republic*
shahlaganbarova@gmail.com

It is shown that when choosing the crystal symmetry, if the wave vector of the wave and the temperature gradient are oriented arbitrarily, waves of a thermomagnetic nature of the same frequency and growth rate are possibly excited.

Keywords: frequency, increment, excitation, symmetry, tensor, orientation.

PACS: 78.55, 73.22.CD, 73.22

INTRODUCTION

It was shown in [1] that hydrodynamic motions in a nonequilibrium plasma, in which there is a constant temperature gradient ($\nabla T = const$), has oscillatory properties. This property of plasma is very different from ordinary plasma. Without an external magnetic field and hydrodynamic motions, transverse “thermomagnetic” waves are possible in it, in which only the magnetic field oscillates. In the presence of an external magnetic field, the wave vector of thermomagnetic waves is perpendicular to the magnetic field and lies in the plane ($\vec{H}, \vec{\nabla}T$). If a weak magnetic field appears in such a plasma, i.e., $\Omega\tau \ll 1$ (Ω - Larmor frequency of electrons, τ - time of collision of electrons) inside the plasma arises in addition to the external electric field, the electric is proportional to the temperature gradient, the electric field is proportional

to the magnetic field. Due to this complex electric field, thermomagnetic waves of a transverse $\vec{k} \perp \vec{\nabla}T$ (\vec{k} - wave vector) and longitudinal character $\vec{k} \parallel \vec{\nabla}T$ are excited. A theoretical study of these thermomagnetic waves in isotropic conducting media of the electric type of charge carriers was carried out in [2-5]. However, in anisotropic conducting media, there is no theoretical study of thermomagnetic waves. In this theoretical work, we will investigate thermomagnetic waves in anisotropic conducting media with selected samples.

BASIC EQUATIONS OF THE PROBLEM

In the presence of an external magnetic field and a temperature gradient in an isotropic solid, the total electric field has the form

$$\vec{E} = \xi \vec{j} + \xi' [\vec{j}\vec{H}] + \xi'' (\vec{j}\vec{H})\vec{H} + A \frac{\partial T}{\partial x} + A' [\vec{\nabla}T\vec{H}] + A'' (\vec{\nabla}T\vec{H})\vec{H} \tag{1}$$

\vec{j} - current flux density

In anisotropic conducting media, all coefficients in equation (1) are tensors.

Then the equations for anisotropic conducting media will have the form:

$$E_i = \xi_{ik} \vec{j}_k + \xi'_{ik} [\vec{j}\vec{H}]_k + \xi''_{ik} (\vec{j}\vec{H}) H_k + A_{ik} \frac{\partial T}{\partial x_k} + A'_{ik} [\vec{\nabla}T\vec{H}]_k + A''_{ik} (\vec{\nabla}T\vec{H}) H_k \tag{2}$$

Here ξ_{ik} is the tensor of the reciprocal of the ohmic resistance, A_{ik} is the differential thermoelectric power, and A'_{ik} is the Nernst-Ettinghausen coefficient. We will consider a solid external magnetic field $\vec{H}_0 = 0$. Then, in the equations, the terms containing are equal to zero. Taking into account Maxwell's equation, we obtain the following system of equations:

$$\begin{cases} E'_i = \xi_{ik} j'_k + A'_{ik} [\nabla TH]_k \\ \text{rot} \vec{E}' = -\frac{1}{c} \frac{\partial \vec{H}'}{\partial t} \\ \text{rot} \vec{H}' = \frac{4\pi}{c} \vec{j}' + \frac{1}{c} \frac{\partial \vec{E}'}{\partial t} \end{cases} \tag{3}$$

Assuming that all variable quantities are monochromatic in nature, i.e.

$$(E', H', j) \sim e^{i(\vec{k}\vec{r} - \omega t)} \tag{4}$$

From (3) it turns out:

$$E'_i = \xi_{ik} j'_k + A'_{ik} [\vec{\nabla}T\vec{H}]_k \tag{5}$$

$$j'_k = \frac{ic^2}{4\pi\omega} \left[\vec{k} \left[\vec{k}\vec{E}' \right] \right]_k + \frac{i\omega}{4\pi} E'_k$$

(ω - oscillation frequency).

THEORY

We (5) it turns out:

$$E'_i = \frac{ic^2}{4\pi\omega} \xi_{ik} (\vec{k}\vec{E}')_k + \frac{i\omega^2 - ic^2k^2}{4\pi\omega} \xi_{ik} E'_k + \frac{cA'_{ik}}{\omega} (\vec{\nabla}TE')_k - \frac{cA'_{ik}}{\omega} (k\vec{\nabla}T)\vec{E}' \quad (6)$$

To obtain the dispersion equation from (6), you first need to choose a coordinate system. We will choose the following coordinate system

$$k_1 \neq 0, k_2 = k_3 = 0, \frac{\partial T}{\partial x_2} \neq 0, \frac{\partial T}{\partial x_3} = 0 \quad (7)$$

Taking into account (7), from (6) it turns out:

$$E'_i = \left(A\xi_{ik}k_e k_k + B\xi_{ik} + \frac{cA'_{ik}}{\omega} k_e \frac{\partial T}{\partial x_k} \right) E'_k \quad (8)$$

When obtaining (8) from (7), we assumed that $\vec{k} \perp \vec{\nabla}T$ i.e. the resulting thermomagnetic waves are transverse.

$$\text{At } E'_i = \delta_{ik} E'_k, \delta_{ik} = \begin{cases} 1, i = k \\ 0, i \neq k \end{cases} \quad (9)$$

From (8) we obtain the following dispersion equations in tensor form

$$N_{ik} = A\xi_{ie}k_e k_k + B\xi_{ik} + \frac{cA'_{ie}}{\omega} k_e \frac{\partial T}{\partial x_k} \quad (10)$$

$$A = \frac{ic^2}{4\pi\omega}, B = \frac{i\omega^2 - ic^2k^2}{4\pi\omega}$$

Expanding by components (10) it turns out:

$$|(N_{ik} - \delta_{ik})| = (N_{11} - 1)(N_{22} - 1)(N_{33} - 1) + N_{31}N_{12}N_{23} + N_{21}N_{32}N_{13} - N_{31}N_{13}(N_{22} - 1) - N_{32}N_{23}(N_{11} - 1) - N_{21}N_{12}(N_{33} - 1) = 0 \quad (11)$$

Here:

$$N_{11} = \frac{i\omega}{4\pi} \xi_{11}, N_{12} = \frac{(i\omega^2 - ic^2k^2)\xi_{12} + 4\pi cA'_{11}k\nabla_2 T}{4\pi\omega}, N_{13} = \frac{i\omega^2 - ic^2k^2}{4\pi\omega} \xi_{13}, N_{21} = \frac{i\omega}{4\pi} \xi_{21},$$

$$N_{22} = \frac{(i\omega^2 - ic^2k^2)\xi_{22} + 4\pi cA'_{21}k\nabla_2 T}{4\pi\omega}, N_{23} = \frac{i\omega^2 - ic^2k^2}{4\pi\omega} \xi_{23}, \quad (12)$$

$$N_{31} = \frac{i\omega}{4\pi} \xi_{31}, N_{32} = \frac{(i\omega^2 - ic^2k^2)\xi_{32} + 4\pi cA'_{31}k\nabla_2 T}{4\pi\omega}, N_{33} = \frac{i\omega^2 - ic^2k^2}{4\pi\omega} \xi_{33}$$

Putting (12) into (11), we obtain the following equation for the oscillation frequency inside an anisotropic body

$$\Phi_5\omega^5 + \Phi_4\omega^4 + \Phi_3\omega^3 + \Phi_2\omega^2 + \Phi_1\omega + \Phi = 0 \quad (13)$$

Solution (13) in a general form is not possible and therefore we will not write out the expression $\Phi, \Phi_1, \Phi_2, \Phi_3, \Phi_4, \Phi_5$.

To solve the dispersion equation (11), we will choose crystals satisfying the following conditions.

- 1) $N_{11} = N_{21} = N_{31}$,
 - 2) $N_{21} = N_{32} = N_{22}$
 - 3) $N_{13} = N_{23} = N_{33}$
- (14)

A crystal satisfying conditions (14) is diagonal.

Taking into account (14), from (11) we easily obtain

$$N_{11} + N_{22} + N_{33} = 1 \quad (15)$$

If

$$\xi_{11} = \xi_{22} = \xi_{33} = \xi \quad (16)$$

from (15) we get

$$\frac{3}{4\pi} \xi \omega^2 - \frac{c^2 k^2}{2\pi} \xi + i\omega'_{21} = 0 \quad (17)$$

$$\omega'_{21} = -cA'_{21}k_1\nabla_2 T$$

From (17) we get:

$$\omega = \omega_0 + i\gamma = \frac{\sqrt{2}}{3} \left(\frac{6\pi|\omega'_{21}|}{\xi} + c^2k^2 \right)^{1/2} + i \frac{\sqrt{2}}{3} \left(\frac{6\pi|\omega'_{21}|}{\xi} - c^2k^2 \right)^{1/2} \quad (18)$$

It can be seen from (18) that at $c^2k^2 > \frac{6\pi\omega'_{21}}{\xi}$, the exciting wave is of a purely electromagnetic nature.

When $c^2k^2\xi < 6\pi|\omega'_{21}|$ the excited wave is growing thermomagnetic with a frequency

$$\omega_0 = \left(\frac{4\pi\omega'_{21}}{\xi} \right)^{1/2} \left(1 + \frac{1}{12\pi} \frac{c^2 k^2 \xi}{|\omega'_{21}|} \right) \text{ and increme } \gamma = \left(\frac{4\pi\omega'_{21}}{\xi} \right)^{1/2} \left(1 - \frac{c^2 k^2 \xi}{12\pi|\omega'_{21}|} \right)$$

For $\vec{k} \parallel \vec{\nabla}T$ from (11), consider the case

$$1) N_{11} = N_{21} = N_{31}, \quad 2) N_{13} = N_{32} = N_{33} = N_{23} = N_{13} \quad (19)$$

$$N_{11} = \frac{i\omega}{4\pi} \xi_{11}, \quad N_{12} = \frac{(i\omega^2 - ic^2k^2) \xi_{12} - 4\pi c \Lambda'_{12} \vec{k} \vec{\nabla}T}{4\pi\omega}, \quad N_{13} = \frac{(i\omega^2 - ic^2k^2) \xi_{13} - 4\pi c \Lambda'_{12} \vec{k} \vec{\nabla}T}{4\pi\omega},$$

$$N_{21} = \frac{i\omega}{4\pi} \xi_{21}, \quad N_{22} = \frac{(i\omega^2 - ic^2k^2) \xi_{22} - 4\pi c \Lambda'_{22} \vec{k} \vec{\nabla}T}{4\pi\omega}, \quad N_{23} = \frac{(i\omega^2 - ic^2k^2) \xi_{23} - 4\pi c \Lambda'_{23} \vec{k} \vec{\nabla}T}{4\pi\omega}, \quad (20)$$

$$N_{31} = \frac{i\omega}{4\pi} \xi_{31}, \quad N_{32} = \frac{(i\omega^2 - ic^2k^2) \xi_{32} - 4\pi c \Lambda'_{32} \vec{k} \vec{\nabla}T}{4\pi\omega}, \quad N_{33} = \frac{(i\omega^2 - ic^2k^2) \xi_{33} - 4\pi c \Lambda'_{33} \vec{k} \vec{\nabla}T}{4\pi\omega}$$

Taking into account (19-20) from (11) we get:

$$N_{11} - 1 + 2N_{11}N_{22} = 0 \quad (21)$$

$$\omega_2 = \frac{i\pi}{\xi} + \left(\frac{2\omega'_{21}\xi}{\pi} \right)^{1/2} \frac{\pi}{\xi} - \frac{i\pi}{\xi} \left(\frac{2\omega'_{21}\xi}{\pi} \right)^{1/2}$$

$$\xi_{11} = \xi_{22} = \xi$$

$$\frac{i\omega\xi}{4\pi} - \frac{\xi^2\omega^2}{8\pi^2} + \frac{\xi^2c^2k^2}{8\pi^2} - \frac{i\omega'_{21}\xi}{2\pi} - 1 \quad (22)$$

From solution (22) we obtain

$$\omega_1 = \frac{i\pi}{\xi} - \left(\frac{2\omega'_{21}\xi}{\pi} \right)^{1/2} \frac{\pi}{\xi} + \frac{i\pi}{\xi} \left(\frac{2\omega'_{21}\xi}{\pi} \right)^{1/2} \quad (23)$$

From (23) it can be seen that a wave with a frequency

$\omega_0 = -\left(\frac{2|\omega'_{21}|}{\pi\xi} \right)^{1/2}$ is increasing, a wave with a frequency

$\omega_0 = -\left(\frac{2|\omega'_{21}|}{\pi\xi} \right)$ can grow if $2|\omega'_{21}|\xi < \frac{\pi}{2}$

For an arbitrary orientation of the wave vector relative to the temperature gradient, from tensor (10), we obtain

$$N_{11} = \frac{i\omega^2 \xi_{11} + \omega_{11}}{4\pi\omega}, \quad N_{12} = \frac{(i\omega^2 - ic^2k^2) \xi_{12} - \omega_{11} + \omega_{12}}{4\pi\omega}, \quad N_{13} = \frac{(i\omega^2 - ic^2k^2) \xi_{13} + \omega_{13}}{4\pi\omega},$$

$$N_{21} = \frac{i\omega^2 \xi_{21} + \omega_{21}}{4\pi\omega}, \quad N_{22} = \frac{(i\omega^2 - ic^2k^2) \xi_{22} + \omega_{22}}{4\pi\omega}, \quad N_{23} = \frac{(i\omega^2 - ic^2k^2) \xi_{23} + \omega_{23}}{4\pi\omega}, \quad (24)$$

$$N_{31} = \frac{i\omega^2 \xi_{31} + \omega_{31}}{4\pi\omega}, \quad N_{32} = \frac{(i\omega^2 - ic^2k^2) \xi_{32} + \omega_{32}}{4\pi\omega}, \quad N_{33} = \frac{(i\omega^2 - ic^2k^2) \xi_{33} + \omega_{33}}{4\pi\omega},$$

$$\omega_{ik} = -4\pi c \Lambda'_{ik} (\vec{k} \vec{\nabla}T)$$

Choosing a crystal.

From (11), taking into account (24), we obtain the following dispersion equation for determining the frequency and growth rate of the excited waves inside an anisotropic crystal

$$\frac{i\omega^2 \xi_{11} - \omega_{11}}{4\pi\omega} \left(\frac{i\omega^2 \xi_{12} - ic^2k^2 \xi_{12} - \omega_{11} - \omega_{12}}{4\pi\omega} + 1 \right) = \frac{i\omega^2 \xi_{22} - ic^2k^2 \xi_{22} + \omega_{22}}{4\pi\omega} \left(\frac{i\omega^2 \xi_{11} - \omega_{11}}{4\pi\omega} - 1 \right) \quad (25)$$

At $\omega_{12} = \omega_{11} + \omega_{22}$ and $\xi_{11} = \xi_{22} = \xi_{12}$ from (25) we get

$$\omega^2 = \frac{1}{2} \left(c^2 k^2 + i \frac{\omega_{12}}{\xi} \right)$$

(26)

Solution (26) gives

$$\omega = \frac{ck}{2} \left(\frac{\omega_{12}}{c^2 k^2 \xi} \right)^{1/2} (1+i) \quad (27)$$

or $\omega = \omega_0 + i\gamma$

$$\omega_0 = \left(\frac{\omega_{12}}{4\xi} \right)^{1/2}, \quad \gamma = \left(\frac{\omega_{12}}{4\xi} \right)^{1/2} \quad (28)$$

For any orientation of the wave vector with respect to the temperature gradient, the frequency and growth rate of the excited thermomagnetic are the same.

DISCUSSION OF THE RESULTS

In anisotropic conducting media of the electric type of charge carriers in an external electric field in the presence of a constant temperature gradient, longitudinal $\vec{k} \parallel \vec{\nabla}T$ and transverse $\vec{k} \perp \vec{\nabla}T$ waves of a thermomagnetic nature are excited. The frequency and growth rate of this wave depend on the conductivity

of the medium. The conductivity of a medium is easily expressed in terms of the diagonal values of the conductivity. This creates a favorable condition for experimental verification of the exciting waves. If the wave vector of the excited waves has an arbitrary direction relative to a constant temperature gradient, then the frequency and growth rate have the same values. When calculating, we choose crystals of different symmetry. Of course, the conditions for the excitation and growth of the wave will be different if we choose different symmetries from the tensor (11).

CONCLUSION

It is proved that in anisotropic conducting media of electric type of charge carriers, different waves of a thermomagnetic nature are excited. With the longitudinal $\vec{k} \parallel \vec{\nabla}T$ and transverse $\vec{k} \perp \vec{\nabla}T$ orientation of the wave vector relative to the temperature gradient, waves of a thermomagnetic nature with different frequencies and increments are excited.

-
- [1] *E.R. Hasanov, A.V. Islamzade, H.Sh. Hasanov.* Thermomagnetic Waves in Anisotropic Conductors International Journal On Technical And Physical Problems Of Engineering, March 2016, Vol.8, №1, pp. 50-54.
- [2] *E.R. Hasanov, I.I. Mustafayeva.* Non-linear fluctuations of concentration of charge carriers and an electric field in semiconductors 13th International Conference on “Technical and Physical Problems of Electrical Engineering”, 21-23 September, 2017, №38, pp. 39-43.
- [3] *E.R. Hasanov, F.F. Aliev.* Nonlinear Oscillation Task for the Concentration of Field Carriers in Semiconductors with Deep Traps IOSR Journal of Applied Physics, Volume 10, Issue 1 Ver. II (Jan.-Feb. 2018), pp. 36-42.
- [4] *E.R. Hasanov, Sh.G. Khalilova, E.O. Mansurova, N.M. Tabatabaei.* Unstable thermomagnetic waves in anisotropic medium of electronic type of charge carriers. International Journal on “Technical and Physical Problems of Engineering” (IJTPE) Turkey, March 2020, Issue 42, Volume 12, №1, pp. 49-52.
- [5] *E.R. Hasanov, Sh.G. Khalilova, Z.A. Tagiyeva, V.M. Hajiyeva, S.S. Ahadova.* Excitation of thermomagnetic and recombination waves in impurity with two types of current carriers International Journal on “Technical and Physical Problems of Engineering”(IJTPE), march 2021, Issue 46, Volume 13, №21, pp. 57-61.

E.R. Həsənov, Ş.G. Xəlilova, G.M. Məmmədova, R.Z. Cavadova

QEYRİ-SABİT ANİZOTROP KEÇİRİCİ MÜHİTLƏRDƏ TERMOMAQNİT DALĞALARI

Göstərilmişdir ki, kristal simmetriya seçildikdə dalğanın dalğa vektoru və temperatur qradienti ixtiyari oriyentasiya edilərsə, eyni tezlikli və artımlı termomaqnit dalğaların həyəcanlanması mümkündür.

Э.Р. Гасанов, Ш.Г. Халилова, Г.М. Мамедова, Р.З. Джавадова

ТЕРМОМАГНИТНЫЕ ВОЛНЫ В НЕУСТОЙЧИВЫХ АНИЗОТРОПНЫХ ПРОВОДЯЩИХ СРЕДАХ

Показано, что при выборе симметрии кристалла, если волновой вектор волны и градиент температуры ориентируются произвольно, возможно возбуждаются волны терромагнитного характера одинаковой частоты и инкремента.