

MONOCHROMATIC WAVE PROPAGATION IN VACUUM

E.R. HASANOV^{1,2}, Sh.G. KHALILOVA², R.K. MUSTAFAYEVA¹¹Baku State University Acad. Z. Khalilov, str.23, Baku, Azerbaijan Republic²Institute of Physics, MSE, AZ-1143, H. Javid 131, Baku, Azerbaijan Republic
shahlaganbarova@gmail.com

Vacuum still remains a mysterious medium from a physical point of view, especially its physical properties. The passing of a monochromatic wave is examined through a vacuum of size L in our work. A monochromatic wave has a certain energy, and when passing through a vacuum it should not lose energy. At different values of the vacuum size, according to the properties of the vacuum, there should be no loss of energy by the wave. If energy loss occurs as a wave passes through, the cause of the energy loss is inhomogeneity. The inhomogeneity of the vacuum proves the existence of some particles in the vacuum, i.e. vacuum is ordinary matter. We can say that a certain part of the energy of a monochromatic wave is absorbed by matter, i.e. vacuum. In this work, the energy of a monochromatic wave is calculated before and after passing through a vacuum of size L . It has been proven that the ratio of the energy of a monochromatic wave before passing through a vacuum is less than 1. This means that the monochromatic wave was scattered inelastically in a vacuum and vacuum is a dense medium. After the passage of the vacuum wave, the length of the monochromatic wave decreases. A monochromatic wave loses energy in a vacuum.

Keywords: vacuum, energy, inelastic interaction, monochromatic wave, absorption, inhomogeneity.

PACS: 78.55, 73.22.CD, 73.22

INTRODUCTION

Quantum mechanics, as a mathematical method, is based on the wave function. In the one-dimensional case, the monochromatic wave function has the form

$$\psi(x, t) = ce^{i(kx - \omega t)} = ce^{i\left(kx - \frac{Et}{\hbar}\right)} \quad (1)$$

Here k - wave vector

$$k = \frac{2\pi}{\lambda} n, (n = 0, \pm 1, \pm 2, \dots), \quad \lambda \text{ - wavelenth, } K \text{ -}$$

energy of wave in vacuum, which is related to the wave vector as follows

$$E = \frac{\hbar^2 k^2 n^2}{2m_0}, \quad n = 0, \pm 1, \pm 2, \dots \quad (2)$$

m_0 - free electron mass

Wave function (1) satisfies Schrödinger equation

$$\frac{\partial \psi}{\partial t} = \frac{i\hbar}{2m_0} \frac{\partial^2 \psi}{\partial x^2} \quad (3)$$

Let's pretend that a monochromatic wave falls on vacuum with energy

$$E_{\text{before}} = \frac{2\pi^2 \hbar^2}{m_0 \lambda_0^2} \quad (4)$$

THEORY

In the view of classical physics, this wave should not change energy after leaving the vacuum with a length of L . However, we will show that the energy (4) changes after leaving the vacuum as L . This means that the vacuum interacts with a monochromatic wave. So, there is interaction with the wave in vacuum. Using the Schrödinger equation (3), we calculate the energy in vacuum

From (1) we calculate $\frac{\partial \psi}{\partial t}$ and $\frac{\partial^2 \psi}{\partial x^2}$

$$\frac{\partial \psi}{\partial t} = 2\pi i \psi \left(-\frac{x}{\lambda^2} \frac{d\lambda}{dt} - \frac{\hbar}{m\lambda^2} + \frac{2\hbar t}{m\lambda^3} \frac{d\lambda}{dt} \right) \quad (5)$$

$$\frac{\partial^2 \psi}{\partial x^2} = -4\pi^2 \psi \left(\frac{1}{\lambda} - \frac{x}{\lambda^2} \frac{d\lambda}{dx} + \frac{2\hbar t}{m\lambda^3} \frac{d\lambda}{dx} \right)^2 + 2\pi i \psi \left[-\frac{2}{\lambda^2} \frac{\partial \lambda}{\partial x} + \frac{2x}{\lambda^3} \left(\frac{d\lambda}{dx} \right)^2 - \frac{6\hbar t}{m\lambda^4} \frac{d\lambda}{dx} + \frac{2\hbar t}{m\lambda^3} \frac{d^2 \lambda}{dx^2} \right] \quad (6)$$

Substituting (5) and (6) into (3) and separating real and imaginary part, we will get

$$2 \frac{\partial \lambda}{\partial x} - \frac{2x}{\lambda} \left(\frac{d\lambda}{dx} \right)^2 + \frac{6\hbar t}{m\lambda^2} \frac{d\lambda}{dx} = 0 \quad (7)$$

MONOCHROMATIC WAVE PROPAGATION IN VACUUM

$$\left(\frac{x}{\lambda^2} - \frac{2\hbar t}{m\lambda^3} \right) \frac{\partial \lambda}{\partial t} = \frac{\hbar}{m\lambda^2} - \frac{2\hbar t}{m\lambda^3} \left(1 - \frac{x}{\lambda} \frac{d\lambda}{dx} + \frac{2\hbar t}{m\lambda^2} \frac{d\lambda}{dx} \right)^2 \quad (8)$$

From (7) we will get

$$\frac{\partial \lambda}{\partial x} = \frac{\lambda \left(2 + \frac{6\hbar t}{m\lambda^2} \right) \hbar}{x} \quad (9)$$

Substituting (9) into (8) we will get:

$$\frac{\partial \lambda}{\partial x} = - \frac{2\pi\hbar}{m\lambda A(t)} \left[1 + \frac{2\lambda}{x} u A(t) + \left(\frac{\lambda}{x} \right)^2 (Au)^2 \right] \quad (10)$$

Here $A(t) = \frac{2\hbar t}{m\lambda^2} - \frac{x}{\lambda}$, $u = 2 + \frac{6\hbar t}{m\lambda^2}$

To integrate equation (10), we assume

$$\frac{2\hbar t}{m\lambda} < x, \quad A = -\frac{x}{\lambda}, \quad u = 2 \quad (11)$$

Substituting (11) into (10) we will get:

$$\frac{\partial \lambda}{\partial t} = \frac{2\pi\hbar}{mx} \quad (12)$$

From

$$\lambda = \lambda_0 + \frac{2\pi\hbar t}{mx} = \lambda_0 \left(1 + \frac{2\pi\hbar t}{mx\lambda_0} \right) \quad (13)$$

When obtaining equation (12), we discarded the member $\frac{\partial^2 \lambda}{\partial x^2}$ and this is quite justified, because, λ depends on the coordinates like a linear way. Thus, the final energy of the wave after leaving the vacuum with size $x = L$:

$$E_{after} = \frac{2\pi^2 \hbar^2}{m_0 \lambda_0^2} \cdot \frac{1}{\left(1 + \frac{2\pi\hbar t}{m_0 \lambda_0 L} \right)^2}$$

That is why

$$\frac{E_{after}}{E_{before}} = \frac{1}{\left(1 + \frac{2\pi\hbar t}{m_0 \lambda_0 L} \right)^2} \ll 1 \quad (14)$$

Expression (14) have gotten from solving equation (10) with values for (11) A and u . For values $A = 0$ and $u = 0$ $\lambda \rightarrow \infty$ and this is the classic area where the wave turns into a line. For values

$$A = \frac{2\hbar t}{m_0 \lambda^2}, \quad u = \frac{6\hbar t}{m_0 \lambda^2} \quad (15)$$

Solution of equation (10) is more complex.

Substituting (15) into (10) for wavelength we will get:

$$\lambda = \lambda_0 \left[1 - 2 \left(\frac{2\hbar\tau}{m_0\lambda_0^2} \right)^2 \left(\frac{3\lambda_0}{L} \right)^2 \right]^{1/6} \quad (16)$$

And

$$\frac{E_{after}}{E_{before}} = \frac{1}{\left[1 - 2 \left(\frac{\hbar\tau}{m_0\lambda_0^2} \right)^2 \left(\frac{3\lambda_0}{L} \right)^2 \right]^{1/6}} \quad (17)$$

Here, τ - the time of passing of wave through vacuum with size L.

Thus, a monochromatic wave, passing through a vacuum of length L, decreases its energy. This is the inelastic interaction of the wave with the vacuum. This means that the vacuum is not an empty means

CONCLUSION

It has been proven that a monochromatic wave, passing through a vacuum of size L, loses part of its energy. Losing energy occurs with different values of L in different ways. Vacuum is matter and has

inhomogeneity. This inhomogeneity leads to losing energy of the monochromatic wave.

A wave with a longer wavelength loses more energy than a wave with a short wavelength. In addition, the wave losing energy leaves the vacuum, which means that the vacuum behaves like a massive medium in which inelastic interaction with the wave occurs. However, unfortunately, it is not known what kind of elementary particles the vacuum is filled with and they move by what law. Vacuum (emptiness) is a mysterious object in the sense of physical understanding. The properties of vacuum are not clearly defined and are closed to people

- [1] *L.D. Landau, E.M. Lifshitz. Quantum Mechanics. Non-Relativistic Theory. Series: Theoretical physics, Moscow: Nauka, 1984, pp.13-37.*
- [2] *L.D. Landau, E.M. Lifshitz. Quantum Mechanics. Non-Relativistic Theory. Series: Theoretical physics, Moscow: Nauka, 1984, pp.42-66.*

- [3] *L.D. Landau, E.M. Lifshitz. Quantum Mechanics. Non-Relativistic Theory. Series: Theoretical physics, Moscow: Nauka, 1984, pp.70-100.*

Received: 08.02.2024